

Annex 4
of Pharmacovigilance Procedure
(subitem 4 of item 2 of section III)

TIME INTERVALS
between use of vaccine, tuberculin and clinical manifestations of adverse reactions by codes

Vaccine, tuberculin	Time intervals between use of vaccine, tuberculin and clinical manifestations of adverse reactions by codes																						
	1 - 2	3.1 - 4.3	5	6 - 8	9	10.1 - 10.4	11	12.1 - 12.2	13	14	15	16.1 - 16.2	17	18.1 - 18.2	19	20	21*	22	23	24	25	26.1 - 26.3	
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
DTP, DT, dT***	<72 hours	<48 hours., up to 7 days**	< 7 days	< 48 hours	-	<72 hours	-	-	-	-	<14 days	<48 hours	<72 hours	-	<48 hours	7 - 14 days	2 - 3 days	-	-	-	-	-	
DTaP, combined vaccines based on DTaP ***	<48 hours	<72 hours, up to 7 days**	<7 days	<48 hours	-	<72 hours	unkn	unkn	-	unkn	<14 days	<48 hours	<48 hours	-	<48 hours	7 - 14 days	2 - 3 days	-	-	-	-	-	
Vaccines for prevention measles, mumps, rubella***	4 - 10 days	24 hours, up to 7 days**	4 - 10 days	4 - 10 days	7 - 10 days	unkn	6 - 15 days	4 - 10 days	7 - 10 days	12 - 25 days	<14 days	<48 hours	<72 hours	-	6 - 12 days	-	-	-	-	-	-	-	-
BCG***	unkn	sr	-	unkn	-	-	-	-	-	-	-	<48 hours	unkn	-	-	-	unkn	1,5 - 6 months	1,5 - 6 months	1 - 12 months	5 - 12 months	1 - 24 months	
OPV***	unkn	-	-	<72 hours	-	<72 hours	-	unkn	-	-	-	-	unkn	4 - 30 days	-	-	unkn	-	-	-	-	-	-
IPV***	<48 hours	<48 hours, up to 7	unkn	<48 hours	-	unkn	-	<48 hours	-	-	<14 days	<48 hours	<72 hours	-	unkn	-	unkn	-	-	-	-	-	-

		days**																				
Hepatitis B vaccine ***	<72 hours	<48 hours, up to 7 days**	unkn	<72 hours	-	<5 days	<72 hours	<72 hours	-	-	<14 days	<48 hours	<72 hours	-	<48 hours	<72 hours	unkn	-	-	-	-	-
Polysaccharide vaccines (Hib)***	<72 hours	<48 hours, up to 7 days**	-	<48 hours	<72 hours	<72 hours	-	<72 hours	-	-	<14 days	<48 hours	<72 hours	-	<48 hours	N/A	2 - 3 days	-	-	-	-	-
Tuberculin***	<48 hours	Pain at injection site or discomfort, irritation or pruritus <72 hours	-	Headache <48 hours	-	-	-	-	-	-	-	-	unkn	-	-	-	-	-	-	-	-	-

* In small premature infants, born at pregnancy week 28 or earlier.

** Only for items 4.1 - 4.3.

*** The list is not comprehensive; it is necessary to use instruction for use of appropriate vaccine and tuberculosis allergen registered in Ukraine

Note. sr – specific reaction to BCG vaccine; unkn – terms of development are unknown.